



INNOVATIVE and AUSTRALIAN

BROADLEAF WEEDER

Selective Turf Herbicide



*Barmac **BROADLEAF WEEDER** is one of the most effective selective broadleaf herbicides available because it incorporates three major, well known weedkillers. Each of the constituents has particular strengths and weaknesses across the wide range of broadleaf weeds which can become a pest in turf situations.*

Active Constituents

325g/L Mecoprop

77g/L MCPA

39g/L Dicamba

By using Barmac **BROADLEAF WEEDER** the strengths of each of the individual weedkillers is picked up on and the weakness overcome. This product is therefore very effective in the control of most problem broadleaf weeds including clover, creeping oxalis and bindii.

BROADLEAF WEEDER cost per hectare is low because of the synergistic action of its components. This means that the effectiveness of the product is more than just the additive effect of its constituents. This also helps to provide a wide range margin of safety to most lawn grasses (**BROADLEAF WEEDER** is not recommended for use on Buffalo grass) as well as non-target plants.



BROADLEAF WEEDER does not contain 2,4-D and therefore is kinder to the environment than some of its competitors. We believe that **BROADLEAF WEEDER** is the best available weed control for golf courses, home lawns, bowling greens, cemeteries, public parks, ovals and turf farms – wherever immaculate turf must be maintained.

Pack Size:

10L

For more information about **BROADLEAF WEEDER** or other BARMAC products visit www.barmac.com.au or one of our distributors.
Barmac Industries Pty Ltd Swanbank Australia

BROADLEAF WEEDER



www.barmac.com.au

DIRECTION FOR USE

| SITUATION | PEST | RATE | CRITICAL COMMENTS |
|---|--|--|---|
| Fairways, Parks, Ovals and playing fields and other turf situations | Bindii, Catsear, Chickweed, Clovers, Cobblers Pegs, Creeping Oxalis (not TAS), | 4.3L in 450L of water per hectare. | Repeat treatments may be necessary and should be made in 7 - 14 days. |
| Bowling & Golf Greens & Lawns | Dandelion, Fat Hen, Fumitory, Lambs Tongue, | 45mL in 20 Litres of water/100m ² . | |
| Newly Sown Turf | Mallow, Pearlwort, Stagger Weed, Wireweed | 30mL in 20L of water per 100m ² | Do not apply until four weeks after the first mowing |