

DIRECTIONS FOR USE:

SITUATION	PEST	STATE	RATE	CRITICAL COMMENTS
<i>Apples</i>	Collar Rot (Sclerotium rolfsii)	Qld, SA and WA	1 kg in 100 L of water.	Drench the seedling butt and planting site with the prepared mixture. Use 500mL of mixture per tree.
<i>Tomatoes</i>	Sclerotium Base Rot (Sclerotium rolfsii)	All States	300-500 g/100 L of water.	Apply 250mL per plant at transplanting applied as a drench. Use the higher rate for heavy soils and high disease levels.
<i>Lettuce</i>	Bottom Rot, Root Rot (Rhizoctonia)	All States	100-150 g/20m ² in 100 litres of water.	Apply when plants are 5-7 cm high and have been thinned. Use the higher rate for heavy soils and high disease levels.
<i>Beans</i>	Stem Rot, Root Rot (Rhizoctonia)	All States	16.5-22 kg/ha in a convenient volume of water.	Apply over row in a band 20-30 cm wide and incorporate well with top 10-15 cm of soil prior to sowing. Use the higher rate for heavy soils and high disease levels.
<i>Peanuts</i>	Crown Rot (Sclerotium rolfsii)	Qld only	16-22 kg/ha.	Spread in 30 cm bands along rows at pegging time. Use the higher rate for heavy soils and high disease levels.
<i>Cotton</i>	Damping Off (Rhizoctonia)	NSW - Qld only	5 kg/ha in a convenient volume of water.	Apply over seed and surrounding soil at planting.
<i>Cotton Seed</i>	Rhizoctonia	NSW and Qld only	240g/100 kg of Cotton Seed.	Apply product as a slurry using a volume of water suited to the seed coating equipment used. Apply to cotton seed in manner which ensures even coverage of the seed. The addition of an appropriate colour dye to indicate pesticide treated seed is a legal requirement in some states and is recommended to prevent its use as a stock feed or for oil production.
<i>Potatoes</i>	Common Scab (Streptomyces scabies)	All States	33-44 kg/ha.	Apply to soil in band 30 cm wide over row at planting time, rotary hoe to 10-15 cm depth. Cover seed with treated soil only. Use the higher rate for heavy soils and high disease levels.
<i>Cabbages, Broccoli, Cauliflowers</i>	Club Root, Wire Stem (Rhizoctonia solani) and Black Rot (Corticium solani)	All States	16.5-22 kg/ha.	Apply as a band treatment over rows and incorporate to a depth of 10 cm prior to planting out. Use the higher rate for heavy soils and high disease levels.
<i>In Seed Bed</i>		All States	500-750 g/100m ² in a convenient volume of water.	Incorporate to a depth of 10 cm. Use the higher rate for heavy soils and high disease levels.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENT: 750 g/kg QUINTOZENE

FOR THE CONTROL OF SOIL-BORNE DISEASES OF VEGETABLES AND ORNAMENTALS.

GROUP Y FUNGICIDE

10 kg
Net Contents



BARMAC INDUSTRIES PTY LTD
Box Flat Estate
Swanbank Road
SWANBANK QLD 4306

APVMA APPROVAL NUMBER 51684/0499

Batch Number:

Date of Manufacture:

SITUATION	PEST	STATE	RATE	CRITICAL COMMENTS
<i>ORNAMENTALS Bulbous Iris</i>	Crown Rot, Black Rot (Sclerotium rolfsii)	All States	150 g/10m ²	Apply evenly and incorporate to a depth of 10-15 cm prior to planting.
<i>Larkspur, Calendula, Sweet-Pea, Snapdragons</i>	Rhizoctonia Stem Rot	All States	120 g/10m ²	Incorporate evenly to a depth of 10-15 cm prior to planting.
<i>Carnations, Begonias, Chrysanthemums, Poinsettias</i>	Rhizoctonia Stem Rot	All States	500-750 g/100m ²	Apply to seed bed either as a drench using 200-500 L of water, or dry, incorporating thoroughly into the top 7-10 cm of soil.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD – PEANUTS AND VEGETABLES: DO NOT APPLY LATER THAN 28 DAYS BEFORE HARVEST.

GENERAL INSTRUCTIONS:

MIXING: Quintozene 750 may be applied to the soil and irrigated then mixed into the top 10-15 cm or mixed with water and applied to the soil surface or prepared as a slurry for seed dressing.

PROTECTION OF LIVESTOCK: Do not feed treated material to animals including poultry.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT: Dangerous to fish. Do not contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL: Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight.

Single rinse plastic bag (where applicable) before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Puncture and bury empty bag in a local authority landfill. If no landfill is available, bury the bag below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty container and product should not be burned.

Triple rinse drums and pails before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return containers to recycler or designated collection point. If not recycling, crush or puncture and bury empty container in a local authority landfill. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty container and product should not be burned.

FUNGICIDE RESISTANCE WARNING: For fungicide resistance management Barmac Quintozene 750 Fungicide is a Group Y Fungicide. Some naturally occurring individual fungi dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Barmac Quintozene 750 Fungicide and other Group Y fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use Barmac Industries Pty Ltd accepts no liability for any losses that may result from the failure of Barmac Quintozene 750 Fungicide to control resistant fungi.

SAFETY DIRECTIONS: Avoid contact with eyes and skin. Wash hands after use.

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre - phone 13 11 26.

MATERIAL SAFETY DATA SHEET: Additional information is listed on the Material Safety Data Sheet available from Barmac Industries Pty Ltd or from our internet site www.barmac.com.au.

Notice: This product is designed only for the particular purposes indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective. The buyer or user is responsible for any misuse or negligence in the handling, storage, use of or disposal of this material. Barmac Industries Pty Ltd will not accept any responsibility whatsoever and howsoever arising whether for consequential loss or otherwise in connection with the supply and use of this product other than those responsibilities mandatorily imposed by Statutes applicable to the sale and supply of these goods.