

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING

**BARMAC**

# SODIUM MOLYBDATE

390g/Kg Molybdenum (Mo) as sodium molybdate

**FOR THE TREATMENT OF MOLYBDENUM  
DEFICIENCY IN VARIOUS SITUATIONS  
AS PER THE DIRECTIONS FOR USE.**

**1Kg**  
Net Weight



**BARMAC INDUSTRIES** PTY LTD

3 Mary Street,  
BLACKSTONE, QLD 4304  
Phone (07) 3280 3000

**Batch  
Number:**  
**Date of  
Manufacture:**

## DIRECTIONS FOR USE

ANNUAL CROPS	g/100L	COMMENTS
General Rate	50-100	Apply in sufficient water to wet foliage. Apply early in crop life.
Seedlings	40	Apply prior to transplanting.
Hydroponics	2.5 ppm can be achieved by mixing 7g of sodium molybdate per 1000L of water	

Note: Molybdenum can be toxic when levels become too high. One spray per crop is generally sufficient, except where deficiency is noted. Susceptible crops such as brassicas and cucurbis may require two sprays three weeks apart.

**PROTECTION OF LIVESTOCK:** Excessive molybdenum can be harmful to stock – do not apply fertilizer containing molybdenum on more than 25% of farm annually. Do not graze for four weeks after application.

*Notice:* This product is designed only for the particular purposes indicated on the label. Failure to use the product strictly as directed may be illegal, prove dangerous and render the product ineffective. The buyer or user is responsible for any misuse or negligence in the handling, storage, use of or disposal of this material. Barmac Industries Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply and use of this product other than those responsibilities mandatorily imposed by Statutes applicable to the sale and supply of these goods.