

CENTER FRONT

11.5"

CENTER BACK

TOMCAT[®]

Rat and Mouse Bait PELLETS



For the control of rats and mice



PELLETS

NET 2.5 kg



DIRECTIONS FOR USE
DO NOT exceed 3m between bait stations for mice, or 9m for rats. DO NOT place bait stations in the open.

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.	Rats (Rattus Norvegicus, Rattus rattus)	150 g per bait tray	Eliminate as far as practicable, all alternative food sources. Bait infested area. Inspect bait stations daily. Replace eaten baits. If eaten quickly increase number of bait stations. Continue observation and replenishment until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.
	Mice (Mus musculus)	150 g per bait tray	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead rodents and unconsumed bait properly. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

PRECAUTION

The use of this product for the control of protected native rodents requires permission from wildlife authorities.

DO NOT allow bait to contaminate foodstuffs or food intended for human or animal consumption.

DO NOT apply bait to crops.

DO NOT place in positions where there is a possibility of contaminating food or surfaces that come into direct contact with food. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces.

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Active Constituent: 0.05 g/kg BROMADIOLONE
Contains Denatonium Benzoate (Bittering Agent)

For the control of rats and mice

including those resistant to other anticoagulants, in and around domestic, commercial, public services and industrial buildings.

NET 2.5 kg

Manufactured by:
Bell Laboratories, Inc.
Madison, WI 53704 U.S.A.
www.belllabs.com

Distributed by:
BARMAC Barmac Pty Ltd
3 Mary Street,
Blackstone, QLD 4304
Phone (07) 3280 3000
www.barmac.com.au

EMERGENCY CONTACT NO. (02) 9984 2255

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT
DO NOT contaminate dams, rivers or streams with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product is hazardous to pigs, cats, dogs, native wildlife and poultry.

STORAGE AND DISPOSAL
Store in the closed, original container in a well ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in locked room or place away from children, animals, food and feedstuffs, seed and fertilizers. Do not dispose of chemicals on site. Break, crush, puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500mm in a disposal pit specially marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burned. Dispose of carcasses safely by burying and burning in an approved landfill.

SAFETY DIRECTIONS
Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. NOT touch bait. When preparing product for use, wear elbow-length PVC gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use wash gloves.

FIRST AID
If poisoning occurs, contact a Doctor or Poison Control Centre. For further information please refer to the Material Safety Data Sheet. Vitamin K (Phytomenadione) is antidotal.

CONDITIONS OF SALE
The use of TOMCAT Rat and Mouse Bait is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories, Inc. regarding its suitability or efficacy for any purpose. Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Trade Practices Act, 1974.

IN AN EMERGENCY DIAL 131 126 POISON INFORMATION CENTRE
NRA APPROVAL NO: 49782 / 01

032504

T801134B SPEC: 23 x 4 (minus 1/2 for gap)

IMAGE



I.T.

DATE	3/26/04
CUSTOMER	Bell Laboratories
ITEM NAME	609PP-2
P.O.#	
IMAGE I.T. JOB#	46113-BLI5-4

PROOF TYPE:
 DIGITAL HP
 DIGITAL KODAK
 PDF
 COLOR KEY
 DYLUX

SIGNATURE OF APPROVAL
1
2
3
THIS PROOF:
 Needs revision
 New proof needed
 Approved with revision
 Is approved